


of 1 molecule of bromide of triethylphosphonium and 1 molecule of triethyl-vinyl-phosphonium,



I have endeavoured to split the latter in accordance with the above equation, but without success.

Triethylphosphine acts with energy upon the homologues of dibromide of ethylene; I have not yet examined, however, any of the products thus obtained. Mr. W. Valentin, to whom I am indebted for much valuable assistance during my experiments, has found, moreover, that triethylarsine unites with dibromide of ethylene. He has not yet completed the investigation of the crystalline body which is generated in this reaction.

II. "On the Different Types in the Microscopic Structure of the Skeleton of Osseous Fishes." By A. KÖLLIKER, Professor of Anatomy and Physiology in the University of Würzburg. Communicated by Dr. SHARPEY, Sec.R.S.
Received January 20, 1859.

After having been occupied for several months with observations on the minute structure of the bones of fishes, I now take the liberty to present the results of my studies to the Royal Society.

The principal fact which I have to mention is, *that a great many genera of osseous fishes possess no bone-corpuscles, radiated or fusiform, in their skeleton, and therefore no real osseous tissue.* That there exist fish-bones without bone-corpuscles must have been long known in England to those who have collections of microscopic preparations of the hard tissues of animals, as Owen, Tomes, Williamson, Quekett, and others; but nobody seems to have mentioned the fact before Williamson, Quekett, Dr. Mettenheimer of Frankfort, and myself*. In the year 1850 Professor Williamson pointed out the absence of bone-corpuscles from the bones of the Cod, Haddock, Perch, Plaice, Pike, and various other fish, distinguishing them in

* Since this communication was read to the Society, Dr. Sharpey has directed my attention to a statement by the late Professor J. Müller, to the effect that in the Pike and many other fish the bones are destitute of bone-corpuscles. This statement occurs in Müller's Annual Report of the progress of Anatomical and Physiological Science in 1835, and is repeated in his addition to the work of Miescher, "De inflammatione Ossium, eorumque Anatome Generali," Berlin, 1836, p. 269.

this respect from the bones of the Eel, in which such corpuscles are abundant*; in 1853 I made known† that the bones of *Leptocephalus* and *Helmichthys* contain no trace of bone-corpuscles; a year later, Mettenheimer showed that the same was true of the bones of *Tetragonurus Cuvieri*‡; and in 1855 Quekett mentions, in the second volume of the 'Histological Catalogue of the College of Surgeons of England,' fishes belonging to eighteen genera, in the bones of which he had not succeeded in finding bone-corpuscles—viz. *Vogmarus islandicus*, *Lophius piscatorius*, *Gadus morrhua*, *Ephippus*, *Sparus*, *Trigla cuculus*, *Belone vulgaris*, *Pleuronectes platessa*, *Trachinus vipera*, *Orthogoriscus mola*, *Exocætus*, *Scarus*, *Esox*, *Sphyræna barracuda*, *Tetrapturus*, *Zeus faber*, *Perca fluviatilis*, *Gobio fluviatilis*. But, notwithstanding these most valuable observations, little or no progress seems to have been made in the more general treatment of this matter, as is best shown by the 'Comparative Histology' of Leydig (1857), in which (p. 157) the *Leptocephalidæ*, *Tetragonurus*, and *Orthogoriscus* are the only cases mentioned, in which the radiated bone-corpuscles are wanting.

On commencing a series of more extended investigations into the minute structure of fish-bones, in October last, I found that the genera which possess real osseous tissue are rather scarce, whilst, on the other hand, I fell in with a great many types in which the bones contained no trace of lacunæ. And as this fact not only appeared to me of interest with regard to the development of the bones of fishes, but also promised to become of great value in systematic zoology, and in the determination of fossil remains, I devoted my whole time to this question. Now that I have investigated more than 200 species belonging to nearly all tribes of osseous fishes, and mounted about 500 microscopic preparations of their hard structures, I hope to be able to treat this question more comprehensively than has been possible hitherto, and in such a way as to lead to some general conclusions.

In giving the results of my observations, I begin with an enumeration of the fishes which belong to the one, and those which belong to the other type.

* Phil. Trans. 1851, p. 693.

† Zeitschr. f. wiss. Zool. iv. p. 361.

‡ Anat.-histol. Untersuch. ü. d. *Tetragonurus Cuvieri*, in den Abh. d. Senkenberg. Gesellschaft, i. p. 241.

I. *Fishes whose bones contain no bone-corpuscles.*

Ordo I. ACANTHOPTERI.

Fam. 1. *Percoidei*.

Pereca fluviatilis.
Apogon Rex mullorum.
Pomatomus telescopium.
Lucioperca sandra.
Serranus cabrilla.
Anthias buphthalmus.
Acerina vulgaris.
Centrarchus sparoides.
Priacanthus macrophthalmus.
Therapon servus.
Trachinus vipera.
Trachinus draco.
Uranoscopus scaber.
Pomotis gibbosus.
Polynemus paradiseus.
Sphyræna spet.
Sphyræna barracuda.
Mullus barbatus.

Fam. 2. *Cataphracti*.

Trigla cuculus.
Trigla lyra.
Prionotus carolinus.
Platycephalus insidiator.
Dactyloptera volitans.
Cottus gobio.
Aspidophorus europæus.
Monocentris japonicus.
Gasterosteus trachurus.

Fam. 3. *Sparoidei* incl. *Mænides*.

Sargus annularis.
Sargus ovis.
Chrysophrys aurata.
Pagrus vulgaris.
Pagellus centrodontus.
Boops salpa.
Boops vulgaris.
Dentex vulgaris.
Smaris vulgaris.
Smaris insidiator.
Gerres Plumieri.

Fam. 4. *Scienoidei*.

Corvina nigra.

Corvina lobata.
Micropogon undulatus.
Otolithus regalis.
Hæmulon formosum.
Pristipoma stridens.

Fam. 5. *Labyrinthiformes*.

Anabas scandens.
Helostoma Temminckii.
Ophicephalus striatus.
Trichopus trichopterus.
Polyacanthus Hasseltii.
Spirobranchus capensis.

Fam. 6. *Mugiloidi*.

Mugil cephalus.
Mugil, spec.
Atherina Humboldtii.
Atherina vulgaris.
Atherina macrophthalma.

Fam. 7. *Notacanthini*.

Mastacembelus pancalus.

Fam. 8. *Scomberoidi*.

Scomber scomber.
Xiphias gladius.
Tetrapturus belone.
Naukrates ductor.
Lampugus pelagicus.
Lampugus sicalus.
Seriola, spec.
Chorinemus saltans.
Caranx trachurus.
Caranx carangus.
Centrolophus pompilus.
Lichia glauca.
Equula insidiatrix.
Argyreus vomer.
Vomer Brownii.
Zeus faber.
Capros aper.
Coryphæna hippurus.
Astrodermus guttatus.
Tetragonurus Cuvieri.

Fam. 9. *Squamipennes*.

Scatophagus argus.

- Holacanthus, spec.
 Toxotes jaculator.
 Ehippus faber.
- Fam. 10. *Tænioidei*.
 Lepidopus argyreus.
 Trichiurus haumela.
 Trachypterus tænia.
 Trachypterus repandus, *Costa*.
 Trachypterus Spinolæ.
 Cepola rubescens.
- Fam. 11. *Gobioidei et Cyclopteri*.
 Gobius capito.
 Gobius cruentatus.
 Gobius longiradiatus, *Risso*.
 Amblyopus Hermannianus.
 Eleotris humeralis.
 Tripauchen vagina.
 Anarrhichas lupus.
 Lepadogaster Gouani.
 Echineis remora.
- Fam. 12. *Blennioidei*.
 Blennius gattorugine.
 Blennius Montaguï.
 Blennius galerita.
 Salaria quadricornis.
 Cristiceps, spec.
 Clinus argenteus.
 Callionymus lacerta.
- Fam. 13. *Pedunculati*.
 Lophius piscatorius.
 Chironectes histrio.
 Maltha vespertilio.
 Batrachus tau.
- Fam. 14. *Theutyes*.
 Naseus longicornis.
 Acanthurus nigricans.
 Amphacanthus javus.
- Fam. 15. *Fistulares*.
 Fistularia tabaccaria.
 Fistularia immaculata.
 Centriscus scolopax.
 Aulostoma sinense.
 Amphisile scutata.

Ordo II. ANACANTHINI, *J. Müll.*

- Fam. 1. *Gadoidei*.
 Gadus æglefinus.
 Gadus morrhua.
 Lota vulgaris.
 Motella tricirrhatta.
 Lepidoleprus trachyrhynchus.
- Fam. 2. *Pleuronectides*.
 Rhombus maximus.
 Rhombus podas.
 Platessa flesus.
 Plaguria, spec.
 Achirus mollis.
- Fam. 3. *Ophidini*.
 Ophidium barbatum.
 Fierasfer imberbis.
 Ammodytes tobianus.
- Fam. 4. *Leptocephalidæ*, *Bp.*
 Helmichthys punctatus.
 Oxystomus hyalinus.
 Leptocephalus pellucidus, *Bp.*
 Hyoprurus messanensis, *Köll.*

Ordo III. PHARYNGOGNATHI, *J. Müll.*

- Fam. 1. *Labroidei cycloidei*.
 Labrus variegatus.
 Labrus serofa.
 Julis vulgaris.
 Julis pavo.
 Crenilabrus pavo.
 Xirichthys novacula.
 Scarus creticus.
- Fam. 2. *Labroidei ctenoidei*.
 Pomacentrus fuscus.
 Dascyllus araucanus.
 Heliases castaneus.
 Glyphisodon rhati.
- Fam. 3. *Chromides*.
 Chromis nilotica.
 Chromis surinamensis.
 Chromis, spec.
 Cichla Deppii.

Fam. 4. *Scomberesoces*.

- Belone vulgaris*.
Belone caudimacula.
Tylosurus imperialis, Bp.
Sayris Camperi.
Hemirhamphus, spec.
Exocoetus exsiliens.

Ordo IV. *PHYSOSTOMI*, J. Müll.Fam. 1. *Siluroidei*.

- Subfam. *Eremophilini*, Bp.
Trichomycterus punctulatus.

Fam. 4. *Cyprinodontes*.

- Pœcilia vivipara*.
Anableps tetrophthalmus.
Cyprinodon calaritanus.
Molienesia latipinnis.
Orestias tæniatus.
Fundulus nigrescens.

Fam. 6. *Esoces*.

- Esox vulgaris*.
Umbra Krameri.

Fam. 7. *Galaxiæ*.

- Galaxias truttaceus*.

Fam. 9. *Scopelini*.

- Saurus lacerta*.
Myctophum elongatum, Bp.
Ichthyococcus Poweriæ, Bp.
Gonostoma denudata, Raf.

Argyropelecus hemigymnus, Cocco.

- Odontostoma Balbo*.

Fam. 10. *Chauliodontidæ*, Bp.

- Chauliodus setinotus*, Schn.
Stomias barbatus, Risso.

Fam. 12. *Heteropygii*.

- Amblyopsis spelæus*.

Fam. 15. *Symbranchii*.

- Symbranchus marmoratus*.
Symbranchus immaculatus.
Amphipnous cuchia.
Monopterus javanicus.

Ordo V. *PLECTOGNATHI*.Fam. 1. *Balistini*.

- Balistes capriscus*.
Monacanthus geographicus.
Aluteres lævis.
Triacanthus brevirostris.

Fam. 2. *Ostraciontes*.

- Ostracion triqueter*.

Fam. 3. *Gymnodontes*.

- Diodon*, spec.
Tetraodon fahaca.
Tetraodon lineatus.
Orthogoriscus mola.

Ordo VI. *LOPHOBRANCHII*.

- Syngnathus typhle*.
Hippocampus guttulatus.
Pegasus draco.

II. *Fishes whose bones contain bone-corpuscles.*Subclassis. I. *Teleostei*, J. Müll.Ordo I. *ACANTHOPTERI*.Fam. 8. *Scomberoides*.

- Thynnus vulgaris*.
Thynnus alalunga.
Auxis bisus.

Ordo IV. *PHYSOSTOMI*.Fam. 1. *Siluroidei*.

- Silurus glanis*.
Silurus bicirrhis.
Schilbe mystus.

- Synodontis serratus*.
Malapterurus electricus.
Malapterurus beninensis.
Heterobranchus anguillaris.
Chaca lophioides.
Plotosus unicolor.
Clarias fuscus.
Pimelodus, spec.
Arius, spec.
Bagrus, spec.
Callichthys, spec.
Loricaria cataphracta.

Auchenipterus furcatus.
Heteropneustes fossilis.
Aspredo laevis.

Fam. 2. *Cyprinoidei.*

Phoxinus laevis.
Cobitis barbatula.
Aspius bipunctatus.
Alburnus lucidus.
Gobio fluviatilis.
Rhodius amarus.
Cyprinus carpio.
Abramis blicca.
Leuciscus rutilus.
Leuciscus tincella.
Tinca chrysitis.
Barbus vulgaris.
Barbus elongatus.
Barbus obtusirostris.
Barbus marginatus.
Chondrostoma risella, Ag.
Dangila lipocheila.
Labeo niloticus.
Catostomus, spec.

Fam. 3. *Characini.*

Citharinus Geoffroyi.
Distichodus niloticus.
Hydrocyon Forskahlii.
Alestes dentex.
Tetragonopterus mexicanus.
Anodus cyprinoides.
Leporinus, spec.
Pacu taeniurus.
Pacu nigricans.
Erythrinus unitaeniatus.
Macrodon trahira.
Piabuca bimaculata.
Gasteropelecus sternicla.
Chirodon, Girard, n. spec.
Brycon, Müll. Tr., n. spec.

Fam. 5. *Mormyri.*

Mormyrus bane.
Mormyrops anguilloides.
Mormyrus longipinnis.
Mormyrus oxyrhynchus.

Mormyrus cyprinoides.
Mormyrus, spec.

Fam. 8. *Salmones.*

Salmo salar.
Salmo trutta.
Argentina silur.

Fam. 11. *Clupeini.*

Clupea harengus.
Alosa vulgaris.
Alosa melanura.
Coilia Grayi.
Engraulis encrasicolus.
Engraulis Brownii.
Notopterus Pallasii.
Macrostoma angustidens, Risso.
Meletta thryssa.
Elops saurus.
Megalops cyprinoides.
Chatoessus cepedianus.
Chatoessus punctatus.
Gnathobolus mucronatus.
Chirocentrus dorab.
Pristigaster, spec.
Lutodeira chanos.
Butirinus macrocephalus.
Hyodon claudulus.
Heterotis niloticus.
Osteoglossum Vandellii.
Osteoglossum formosum.
Sudis gigas.
Alepocephalus rostratus.

Fam. 13. *Muraenoidi.*

Anguilla vulgaris.
Conger myrus.
Ophisurus serpens.
Nettastoma melanura.
Sphagebranchus imberbis.

Fam. 14. *Gymnotini.*

Gymnotus electricus.
Carapus brachyurus.

Subclassis II. *Ganoidei.*

Ordo I. *HOLOSTEI.*

Fam. 1. *Lepidosteini.*

Lepidosteus platyrhynchus.

Fam. 2. *Polypterini*.
Polypterus bichir.
 Fam. 3. *Amiidae*.
Amia calva.
 Ordo II. CHONDROSTEI.
 Fam. 1. *Acipenserini*.
Acipenser naccarii.

Scaphyrhynchus Rafinesquii.
 Fam. 2. *Spatulariæ*.
Spatularia folium.
 Subclassis III. *Dipnoi*.
 Fam. 1. *Sirenoidei*.
Lepidosiren annectens.

From these facts it follows that the osseous fishes, notwithstanding their great number, are separated in a very remarkable way into two groups, as shown in the following enumeration :—

Fishes with bone-corpuscles.

- I. All the extensive and higher-organized tribes of *Physostomi*, J. Müll. ; viz. the
 - Siluroidei (except *Trichomycterus*).
 - Cyprinoidei.
 - Characini.
 - Mormyri.
 - Salmones.
 - Clupeini.
 - Muraenoidei.
 - Gymnotini.
- II. All the *Ganoidei*.
- III. The *Sirenoidei*.
- IV. Of the *Acanthopteri*, only the genus *Thynnus*, Cuv.

Fishes without bone-corpuscles.

- I. All the numerous tribes of the *Acanthopteri*, with the exception of the genus *Thynnus*.
- II. All the *Anacanthini*, J. Müll.
- III. The *Pharyngognathi*, J. Müll.
- IV. Some smaller and lower-organized tribes of *Physostomi*, as the
 - Cyprinodontes.
 - Esoces.
 - Galaxiæ.
 - Scopelini.
 - Chauliodontida, Bp.
 - Heteropygii.
 - Symbranchii.
- And of the Siluroids, only the genus *Trichomycterus*.
- V. The *Plectognathi*.
- VI. The *Lophobranchii*.

As there can be no doubt that most of the higher-organized fishes are amongst those with bone-corpuscles, and as we know that amongst the higher vertebrata, even the lowest, viz. the Perennibranchiata, possess real osseous tissue, it seems to follow that the peculiar distribution of real osseous tissue and of the "osteoid" structure, as the osseous tissue without corpuscles may be called, has a deeper signification. This will be found by studying the development of the bones in both groups ; and I hope to be able, before long, to present to the Royal Society some new facts with regard to this matter also ; but in the mean time, until my observations are more complete, I must abstain from further explanation.

The facts exposed hitherto have had reference only to a great and fundamental structural difference between two extensive groups of osseous fishes. I may now add, that there exist also greater or lesser structural discrepancies amongst the different tribes of each group. But as this is not a suitable occasion for an exposition of the details of this question, I will only say this much :—In the higher fishes, those with real osseous tissue, there exist differences, especially with regard to the *form* and *size* of the bone-corpuscles ; and I hope to be able to show that there are peculiar and tolerably well characterized types of them amongst the Ganoids, Siluroids, *Salmonidæ*, Cyprinoids, *Clupeini*, &c. In the second group there are more varieties. In some tribes the bones are quite structureless homogeneous masses, as in the *Leptocephalidæ* ; in others they have a peculiar fibrous appearance, and consist of a singular mixture of cartilage and osteoid structures, as Quekett first showed in the genera *Orthogoriscus* and *Lophius*, to which I may add some *Balistini* ; but in the great majority of the tribes of this group, the bones contain peculiar *tubes* more or less similar to those of dentine. If these tubes are well developed, the bones acquire a structure *which can in no way be distinguished from that of dentine*,—a fact, which also did not escape the perspicacity of Quekett, who mentions its occurrence in the genus *Fistularia*, the Barracuda Pike (*Sphyræna barracuda*), and the Gar-fish (*Belone vulgaris*). I found the same structure in many other genera of this group, especially among the *Plectognathi*, *Pharyngognathi*, *Sparidæ*, and *Squamipennes* ; but in the greater number this tubular structure is not so well developed, and is intermingled with more structureless parts. Another fact deserving of mention with regard to the bones of this group is, that there very frequently occur also structures, formed by the agglomeration of calcareous globules of different sizes, which resemble in a remarkable degree the lower layers of common fish-scales.

My observations have also extended to the *hard structures of the skin* of fishes, and of *the rays of the fins* ; and I may say that in general the same laws, which apply to the structure of the endoskeleton, hold good also for the exoskeleton. Evidence of this is especially afforded by the fins, the rays of which, independently of their hard or soft condition, contain bone-corpuscles in all the tribes where the internal bones are provided with them, whilst in the other

case these rays are formed of a homogeneous osteoid substance or of a tubular structure, which may also in some fishes, as Williamson first showed in the Ostracions, assume the structure of real dentine, as in many Plectognaths (*Triacanthus*, *Monacanthus*, *Aluterus*, *Tetraodon*, and others), and in certain *Acanthopterygii* (*Equula*, *Ephippus*, *Hæmulon*, *Pristipoma*, *Scatophagus*, *Centrarchus*). With regard to the skin, we may at least go so far as to say that in no fish whose endoskeleton is destitute of bone-corpuscles do they exist in the hard structures of the skin; but, on the other hand, the tribes which have real osseous tissue do not all present it also in the skin. Scales or plates with bone-corpuscles are found amongst living Ganoids, *e. g.* in *Polypterus*, *Lepidosteus*, and even *Amia* (in whose scales J. Müller erroneously supposed them to be wanting), and also in the *Acipenserini* and *Spatulariæ*; they exist also in the fossil Ganoids, as the excellent observations of Williamson have shown.

In many Ganoids, moreover, as Williamson and Quekett have shown, the scales often contain dentinal tubes, or even portions of real dentine ("Kosmine" of Williamson) amidst true bone. In the scales of *Lepidosiren*, also, I find bone-corpuscles, but mostly fusiform, and only here and there having a simple stellate figure. Of the other fishes which have bone-corpuscles in their skeleton, little has hitherto been noted as to the coexistence of such corpuscles in their scales, but I find it to prevail to a considerable extent among them. The presence of bone-corpuscles has been long known, it is true, in the larger scales of the "corselet" of *Thynnus*, also in the dermal plates of certain Siluroids (*Loricaria* and *Callichthys*), and was pointed out by J. Müller in the scales of *Sudis*. Leydig, too, states that true bone-corpuscles exist in the walls of the grooves and semicanals upon the scales of the lateral line in certain Cyprinoids (Carp, Tench, and Barbel). This statement I am able fully to confirm, and to add the following genera in which I find the same thing to occur; viz.—*Hydrocyon*, *Alepocephalus*, *Macrostoma*, Risso, *Pia-buca*, *Serrasalmo*, *Xiphorhamphus*, *Tetragonurus*, *Salminus*, *Chalcinus*, *Pygocentrus*, *Labeo*, and *Catostomus*. But, besides the instance of *Sudis* and certain Siluroids above referred to, I find that many other *Physostomi* have true bone-corpuscles in their scales; not only those of the lateral line, but all of them. From the results of my examinations up to this time, which, however, on account of

the want of materials, are by no means complete, I am able to make out the following list :—

I. CHARACINI.

Of this family I have had the means of examining nearly all the genera, including forty-one species.

Characini with bone-corpuscles in all their scales.

Erythrinus unitæniatus, <i>Spix.</i>	Anodus cyprinoides, <i>Müll. Tr.</i>
Erythrinus microcephalus, <i>Agass.</i>	Anodus edentulus, <i>Agass.</i>
Macrodon trahira, <i>J. Müll.</i>	Anodus leucos, <i>de Fil.</i>
Macrodon auritus, <i>Val.</i>	Schizodon fasciatus, <i>Agass.</i>
Pacu tæniurus (Prochilodus tæniurus, <i>Val.</i>).	Chilodus punctatus, <i>Müll. Tr.</i>
Pacu nigricans, <i>Spix.</i>	Rhaphiodon (Cynodon) vulpinus, <i>Agass.</i>
Pacu lineatus, <i>Val.</i>	Leporinus fasciatus, <i>Müll. Tr.</i>
Distichodus niloticus, <i>Müll. Tr.</i>	Leporinus elongatus, <i>Val.</i>
Alestes dentex, <i>Müll. Tr.</i>	Citharinus latus, <i>Ehr.</i>

Characini without bone-corpuscles in their scales.

*Hydrocion Forskahlîi, <i>Cuv.</i>	Myletes rhomboidalis, <i>Cuv.</i>
*Piabuca bimaculata (<i>Hyrtl. misit</i>).	Tetragonurus mexicanus, <i>de Fil.</i>
Gasteropelecus sternicla, <i>Bl.</i>	*Tetragonurus argenteus, <i>Art.</i>
Gasteropelecus securis, <i>de Fil.</i>	*Tetragonurus maculatus, <i>Müll. Tr.</i>
Cheirodon Girard, nov. sp., <i>de Fil.</i>	*Salminus Orbignyanus, <i>Val.</i>
Brycon falcatus, <i>Müll. Tr.</i>	*Chalcinus Mülleri, <i>de Fil.</i>
Brycon, nov. sp., <i>de Fil.</i>	Pygocentrus nigricans, <i>Müll. Tr.</i>
Serrasalmo rhombeus, <i>Cuv.</i>	Epiplatys gibbosus, <i>Müll. Tr.</i>
*Serrasalmo marginatus, <i>Val.</i>	Piabucina erythrinoides, <i>Val.</i>
Xiphorhamphus falcatus, <i>Müll. Tr.</i>	Exodon paradoxus, <i>Müll. Tr.</i>
*Xiphorhamphus hepsetus, <i>Müll. Tr.</i>	Leporinus, spec.
Myletes rubripinnis, <i>Müll. Tr.</i>	

With regard to the members of the second division, it is to be observed, that probably in all of them the canals attached to the scales of the lateral line are formed of true osseous tissue ; in those marked with an asterisk I have found this by actual examination.

The *Characini* are thus divisible into two groups, according to the nature of their scales ; at the same time, these are not to be regarded as natural divisions in other respects, and the less so as one and the same genus, such as *Leporinus*, for example, may include species which differ in the composition of their scales. The presence of corpuscles, though connected partly with the size of the scales, does not depend solely on this, for they may be wanting in large scales

(*Hydrocyon*, *Chalcinus*, *Salminus*), and present in small ones (*Anodus edentulus*, *Chilodus*).

2. MORMYRI.

Mormyrus longipennis, Rüpp.	Mormyrus cyprinoides.
Mormyrus oxyrhynchus.	Mormyrus, spec.
Mormyrus bane.	Mormyrops anguillaris.

3. CLUPEINI.

Megalops cyprinoides.	Butirinus macrocephalus.
Elops saurus.	Hyodon claudulus.
Coilia Grayi.	Osteoglossum Vandellii.
Notopterus Pallasii (corpusc. very scanty).	Osteoglossum bicirrosum.
	Heterotis niloticus.

The plates of the abdominal carina in many *Clupeini* are formed throughout of true bone, but do not belong to the present category.

I am unable to find corpuscles in the scales of *Lutodeira chanos*, *Chatoessus punctatus* and *cepedianus*, and *Alosa vulgaris*. In several Cyprinoids (*Labeo*, *Catostomus*, *Barbus*), I have, in like manner, failed to discover corpuscles in the scales proper; on the other hand, I have found very distinct dentinal tubes in the scales of *Barbus*, at their hinder part.

True osseous tissue will doubtless hereafter be found in the scales of many other *Physostomi* which have it in their skeleton, but it is not to be supposed that this will apply to all.

In the *Physostomi*, as in the Ganoids, the bone-corpuscles lie in the lower stratum of the scale; still they are situated above the fibrous layer, and immediately beneath the structureless layer, to which in all scales I apply the name of "ganoin-layer," inasmuch as it has in all cases the same signification.

From the foregoing observations we are able to show still more positively than could be done by J. Müller, that the scales of Ganoids have no peculiarity of structure to distinguish them from those of the *Teleostei*. Nay, certain Ganoids, as *Amia*, have scales, which in respect even of pliancy, rounded contour, and the surface-marking of the ganoin-layer, agree with those of other fishes.

In reference to those fishes which want bone-corpuscles in their skeleton, I have still to remark,—1, that the corpuscles are also invariably wanting in the semicanals upon the scales of the lateral line; for what Leydig designates as rudimentary bone-corpuscles in the

Perch are in fact the tubules of the osteoid substance ; 2, that amongst the group of fishes in question, there are some which have beautiful dentine in their skin-bones, *e. g.* *Amphisile scutata* and the Ostracions.

To the foregoing remarks on the microscopic structure of the hard tissues of fishes, I may add, that there also exists a third group of fishes, in which the endoskeleton is composed only of common cartilage, or of cartilage with depositions of earthy salts, viz. the *Cyclostomi* and *Selachii*. None of these fishes, not even the *Plagiostomi* and *Chimæra*, possess real bone-cells in their hard parts ; for these are formed only, as J. Müller showed many years ago, by ossified cartilage, that is to say, cartilage-cells in an ossified matrix. Even the hard spines of the fins and of the skin of these animals are not real bone, but dentine, as was demonstrated long since by Agassiz and Quekett.

If now we sum up all that has been said, we arrive at the following conclusions :—

I. There exist *three types* of structure in the skeleton of fishes, viz. :

1. *Type of the Selachii.*

The skeleton is formed of cartilage or ossified cartilage.

Selachii, Cyclostomi.

2. *Type of the Acanthopterygii.*

The skeleton is formed of a homogeneous or tubular osteoid substance, often of true dentine.

Teleostei (*J. Müll.*), with the exception of the greater part of the Physostomi (*J. Müll.*).

3. *Type of the Ganoidei.*

The skeleton is formed of real osseous tissue.

Most of the Physostomi, the Ganoidei, and Sirenoidei.

II. The *exoskeleton* follows in some respects the same laws as the endoskeleton, and shows the following types :—

1. *Exoskeleton formed of a homogeneous and fibrous osteoid substance.*

Scales of the majority of the Teleostei.

2. *Exoskeleton formed of dentine.*

Spines of Selachii and scales of Plectognathi, and of *Amphisile*, in part.

3. *Exoskeleton formed of real bone* ; partly in association with homogeneous osteoid substance (*ganoïn*) and dentinal tubes.

Scales of Ganoidei, of *Lepidosiren*, some Siluroidei, of Mormyri, many Characini and Clupeini, also of *Thynnus*.

In terminating this communication, I think it proper to mention that the great liberality with which my friend Mr. Tomes of London, and Professor Williamson of Manchester, put their large collections of microscopic preparations of teeth, bones, and scales at my disposal, proved of great assistance in my investigations, and, accordingly, I am only fulfilling an agreeable duty in now publicly expressing my obligations to them. I am also greatly indebted to my friends Filippo de Filippi of Turin and Henry Müller of Würzburg, also to Dr. Hyrtl of Vienna, and Dr. Peters of Berlin, who supplied me with many of the rarer Mediterranean and foreign fishes. But, in order that my observations may yield the results which may not unreasonably be expected from them, I need more aid ; and as England is the country in which not only the largest zoological collections of fishes, but also the greatest number of microscopic preparations of the hard tissues of recent and fossil animals, are to be found, I take the liberty to ask the possessors of such collections who may be interested in this matter to favour me with such specimens as may seem to them calculated to give to this series of observations the greatest possible extension.

III. "On the Physical Phenomena of Glaciers."—Part II. By Dr. TYNDALL, F.R.S. Received February 24, 1859.

[Abstract.]

The main portion of this Paper deals with the *veined structure* of glacier ice. The author refers to his observations in the Mer-de-glace in 1857, and his reasons for withholding them, and visiting the glaciers once more in 1858.

He describes the general aspect of the structure, and examines the two theories of the phenomenon which are now deserving of attention ; one of these considers the blue veins to be a continuation of the bed-